



## Paediatric Cataract-Type of Surgery Done in my Current Sample of Cases

Authors

**Dr N. Suhasini M.S. FIGS<sup>1</sup>, Dr V. Suryanarayana M.S.,D.O<sup>2</sup>.**

<sup>1</sup>Asst. Prof-KIMS & RF, Amalapuram, <sup>2</sup>Retd. Prof, REH, Visakhapatnam

### Abstract

**Aim:** To describe and analyze type of surgery done in the management of paediatric cataracts.

**Materials & Methods:** A retrospective study of 68 subjects who had undergone cataract surgery during 2007-08 in Govt. REH. PMMA Lens (12mm) was implanted in the bag for children more than 2 years. Intraocular lenses were not implanted for children aged less than 2 years. Optical correction was given.

**Results & Discussion:** SICS+PCIOL done in 29 subjects, phaco + PCIOL done in 15 subjects, only ECCE in 2 patients, ECCE+ACIOL in 1 patient & ECCE + PPC & PCIOL in 1 patient.

### Conclusions:

1. Improvement of child health care is needed for early detection of cataract in children.
2. Role of rubella and trauma in childhood cataract should be investigated and addressed. Visual assessment and postoperative care should be further improved.
3. Cataract surgery with unilateral and bilateral IOL implantation can provide a beneficial effect on final visual outcome in children who are operated- on, before abnormal foveolar function develops.
4. Primary posterior chamber lenses are recommended for children six years of age and older
5. Proper case selection and a controlled, skilled surgical approach, the use of an intraocular lens for visual rehabilitation in the pediatric age group is a feasible approach.
6. Intraocular lens implantation for children with congenital or developing cataract is an effective treatment for visual rehabilitation, even for those patients age 2 years and younger.
7. If nystagmus has developed, the amblyopia is unfortunately irreversible.
8. Concerted efforts must be made to educate people about the prevention of blindness due to cataract in all age groups in general and the importance of continued follow up in the paediatric age group in particular.
9. Apart from school screening programmes, training programmes must be conducted to equip teachers to identify vision problems and to sensitise them about the significance of early reporting.
10. The facilities at primary Health care institutes need to be upgraded and the staff trained to provide postoperative follow up care to paediatric patients.
11. Although the treatment of childhood cataract has evolved considerably over the years, providing high quality standard eye care still remains a challenge for the providers
12. The eye care providers should collaborate with the programs related to child health care and address the underlying causes of cataract in children
13. For this, the cooperation of parents, paediatricians & ophthalmologists is required.

**Keywords:** PMMA-Poly Methyl Methacrylate; REH: Regional Eye Hospital.

### **Management of paediatric cataract**

The management of cataract in the child depends upon several factors: the child's age; the extent of visual loss; whether or not there is an associated illness or coexisting eye disease; and if the cataract Affects One or both eyes.

### **Surgery**

Intraocular lens (IOL) implantation is becoming an increasingly accepted procedure in young children and infants. Awareness of the rate of myopic shift which takes place in the developing eye, and the use of biometry helps to predict appropriate IOL powers to try to ensure eventual emmetropia. However, final refraction is variable, such that emmetropia in adulthood cannot be guaranteed, as there are insufficient long-term studies. Whether or not an IOL is inserted will depend on the age of the child (recently in some centre IOLs are inserted into the eyes of babies as young as four weeks of age), whether or not there is any underlying ocular pathology in which IOL implantation is contra-indicated. In some instances, for example, in remote areas in less industrialised countries where it is unlikely that the child will be able to be brought back for regular follow-up, IOL implantation is not advisable

### **Bilateral Cataracts**

Surgery is indicated when the cataracts are interfering with normal visual development. If the vision is good, then surgery is not indicated, and the child has regular checkups. If there is a significant deterioration in that child's acuity at any stage, then surgery is considered .If a child has bilateral dense cataracts, one eye is operated on followed by the second eye one to two weeks later. The first eye is occluded post-operatively until the second eye is operated on, to prevent amblyopia in the un-operated eye. In very young babies, an IOL is not inserted; these babies are given contact lenses one to two weeks post operatively, and if contact lenses are contra-indicated, the child is given aphakicspectacles. In some of these children, an IOL may be inserted as a secondary procedure when the eye has grown. In

general, an IOL is inserted into the eye of older babies, and depending on the age of the child, an IOL power is selected (according to the biometry) to make that eye appropriately hypermetropic, and hence allow for the myopic shift which takes place in the developing eye; the aim is for emmetropia when the eye is fully grown. In the older child, for example a seven year old child with bilateral cataracts which are beginning to significantly interfere with visual acuity, the eye with the worst visual acuity is operated on, followed by the second eye at least three to four weeks later. All surgery has potential complications, and the parents are warned about all possible complications pre-operatively (see later). Complications of lens aspiration and IOL implantation include post-operative uveitis, endophthalmitis, glaucoma and retinal detachment. In addition, there is a greater than 90% risk that the posterior capsule will become thickened within in a year, which requires a further procedure either with laser or surgical capsulotomy to clear the visual axis and restore visual acuity to what it was initially postoperatively. In order to reduce the need for repeated general anaesthetics in order to perform laser and or surgery, a primary posterior capsulorrhexis (and limited anterior vitrectomy) may be performed. This involves making a hole in the posterior capsule and removing a small amount of vitreous at the time of the initial cataract surgery. This procedure significantly reduces the development of posterior capsular opacification.

### **Unilateral Cataract**

In infants with unilateral cataract, the cataract is most likely to be idiopathic and there is usually no associated systemic disease. It is important not to confuse asymmetric bilateral cataract with unilateral cataract and, hence, careful slit lamp examination is essential in all children with cataract, as there may be very subtle changes in one lens and dense cataract in the other. With regard to the management of unilateral cataract, the decision to operate depends on the density of

the lens opacity, whether or not there is any associated microphthalmos, and on the parent's motivation to have the surgery performed.

The parents need to be fully informed that surgery alone will not improve their child's vision, and that the management of amblyopia is the key to a successful result. This entails optical correction and many hours of patching the normal eye every day, which as one knows can be extremely difficult in a small infant or toddler, and needs to be maintained for many years. If occlusion is not performed the eye will remain densely amblyopic. It has been reported extensively in the literature that children who have had surgery for bilateral cataracts have a much better visual outcome than children who have had surgery for unilateral cataract. Children with unilateral cataract need diligent occlusion therapy of the unoperated eye together with accurate optical correction of the operated eye to obtain a good visual result.

### **Optical correction post-operatively**

#### **Aphakia**

If the child has not had an IOL inserted and is therefore aphakic, contact lenses are fitted as soon as possible after surgery, regardless of age. This is usually done one to two weeks post-operatively. The baby is refracted and contact lenses are fitted. Contact lenses are relatively safe (Amaya, L., et al., 1990), they can be used in combination with spectacles for near, and also to correct aniseikonia (Enoch, J.M. and Hamer, R.D.,1983). Rigid gas permeable contact lenses are most suitable for the management of aphakia, as they have a wide range of powers, can correct astigmatism, and are relatively cheap. The disadvantages of contact lens wear are similar to those seen in adults, and include hypoxia and infective keratitis. Parents should be educated about contact lens hygiene and the signs and symptoms of possible complications associated with contact lens wear. All infants and children wearing contact lenses should be kept under constant review at a specialized unit. If contact lenses cannot be fitted (e.g. raised intraocular pressure, external eye disease, dry eye,

severe microphthalmos or parents who are unable to cope with contacts), the child is given aphakic spectacles which have the advantage of increasing magnification and, therefore, enhancing the child's acuity, and in can make the eyes of children with microphthalmia look slightly bigger.

#### **Pseudophakia**

To allow for the future growth of the infant eye (i.e the myopic shift) an IOL power less than that to result in emmetropia is used. This makes the child hypermetropic (the amount of hypermetropia depends on the age of the child), then as the child's eye grows emmetropia is approached. The child is then given reading spectacles (and a distance prescription if required).

#### **Amblyopia management post-operatively**

The treatment of amblyopia is essential in order to obtain a successful outcome in patients who have had both unilateral and bilateral cataract surgery. Amblyopia is managed by giving the required optical correction and occluding the eye with better visual acuity. The occlusion is performed according to a standardized regime (Tayloe, D.S.I., The Doyne Lecture,1998).It is important to realize that excessive patching of the phakic eye in infants with unilateral aphakia may be associated with increased nystagmus and can have adverse effects on the phakic eye, hence it is imperative that these children are reviewed every two weeks.

#### **Complications of Cataract Surgery**

Cataract surgery in the child and especially in babies has a higher incidence of complications than those seen after adult cataract surgery. This may be due to several reasons: cataract surgery is technically more difficult in the child's eye, as the tissues are more elastic and behave differently to adult tissues; the child's eye becomes more inflamed than an adult eye in response to an IOL; the very young child's visual system is still developing and is therefore extremely sensitive to having a defocused image as is the case in aphakia; and a young child is less likely to fully comprehend the importance of keeping dirty fingers away from the operated eye in the immediate post-operative period and, therefore,

may rub the eye too vigorously resulting in wound dehiscence and iris prolapse or infection. There have been many reports of the visual outcome and complications of posterior chamber lens implantation in children (Thouvenin, D., 1995).

The most common complications include:

### 1. Glaucoma

Glaucoma may arise in the first few weeks post-operatively, and may present as a watery eye with or without photophobia. The child is usually irritated or just "not his usual self". Some children are asymptomatic, hence the importance of regular review with IOP checks post-operatively. Glaucoma may also occur as a late complication years later, and this type of glaucoma can be asymptomatic, therefore all children who have had cataract surgery should be followed up for life.

### 2. Uveitis

All children will develop a certain degree of uveitis post-cataract surgery. This is kept under control with frequent installation of topical steroids, which are given to all children routinely, and are eventually tailed off and discontinued. However, some children (approximately 30%) develop inflammation with fibrin formation. This accumulation of fibrin may result in secondary membrane formation blocking the visual axis if not recognized and treated promptly.

### 3. Irregular pupil

An irregularly shaped pupil after cataract surgery may be due to: a) a strand of vitreous coming forwards from the vitreous cavity to the anterior chamber; b) damage to the iris during surgery; and c) iris prolapse, which may be due to accidental trauma to the eye post-operatively and requires urgent attention, as if left untreated that eye may become infected, and even the other eye may be at a risk from sympathetic ophthalmia.

### 4. Endophthalmitis

Bacterial infection of the eye is a devastating complication which can occur in up to 0.4% of eyes after cataract surgery. It may occur early (i.e. within days to weeks) or late (months to years) post-operatively and presents with a sore painful

red eye, and/or reduced vision, and requires urgent hospital admission and treatment.

### 5. Posterior capsule opacification

Posterior capsule opacification occurs in up to 90% of paediatric eyes after lens aspiration when the posterior capsule and anterior vitreous face are left intact. These children, having had a good visual result initially, later present with blurred vision. The thickened posterior capsule is seen using the slit lamp, or simply by examining the red reflex. These children are admitted as a day case and a posterior capsulotomy is performed. Older children can have laser treatment without a general anaesthetic. In some cases, the capsule thickens again and may require surgery to clear the visual axis.

### 6. Retinal detachment

Retinal detachment may occur years after cataract surgery, and present with reduced vision which may be preceded by flashes and floaters.

## Methodology

### Background

Early diagnosis of congenital cataract is essential in ensuring treatment, together with parental advice and support, can be provided promptly. There is good scope for investigating them using many approaches. Many questions can be easily addressed through trials and clinical studies. Visual loss in congenital/infantile cataract is mainly due to amblyopia. There is limited information available about the long-term outcomes in children treated with modern surgical techniques. This study is designed to evaluate Paediatric cataracts and its visual outcome by using modern techniques among outpatient population of ophthalmology in and around Guntur Township of Andhra Pradesh.

### Objectives

1. To determine functional results after unilateral and bilateral cataract surgery in children with different aphakic optical correction.

2. To report treatment of pediatric patients with cataract, and evaluate the efficiency of different surgical interventions.

The specific objectives are:

To examine does cataract extraction combined with primary IOL implantation, compared with cataract extraction without IOL implantation, result in:

- Better overall visual function?
- Reduced risk of aphakic glaucoma?
- Increased risk of significant myopic shift?

Random sampling is used to select a nationally representative sample of adults. The study design and protocol are submitted to NTR University of Health Sciences for approval.

Each subject and his parent was interviewed, a brief history of the age of onset of visual loss, involvement of other members of the family, the place of residence, and whether the parents' marriage was consanguineous was recorded. Every subject had their visual acuity measured and underwent auto refraction and fundus/optic disc examination.

An exploration of demographic variables was conducted using regression modeling. For the study, 60 subjects of cataract, 0- 18 years age groups underwent an extensive ophthalmologic screening examination, including measurements of visual acuity and the visual field and fundus photography. Cause of cataract formation in each subject was investigated using all screening information and medical records, to establish the aetiology.

### Surgical Technique

Wound construction was done using self-sealing suture less wound in most cases. Capsular

management was obtained in most cases through a central capsulorrhexis measuring around 4 mm. Lensectomy and Anterior vitrectomy were done in children under 2 years and in traumatic cases where Posterior capsule was suspected to be ruptured, a limbal approach was taken for lensectomy and anterior vitrectomy. A primary posterior capsulotomy with or without vitrectomy was performed. PMMA lens (12 mm) was implanted in the bag for children above 2 years. Intraocular lenses were not implanted for children aged less than 2 years. Children who did not receive IOL implants were rehabilitated postoperatively using spectacles or contact lenses. Patching was done if the cataract was unilateral and amblyopia was present.

The surgical procedures included cataract extraction, intraocular lens implantation, posterior capsulorrhexis and anterior vitrectomy in most of the cases. The visual status of eyes with cataract before and 6 weeks after surgery has been evaluated.

### Statistical analysis

The results thus obtained are analyzed, and discussed in comparison with the existing studies in the literature. SPSS Version 10.1 software has been used for statistical analysis. Descriptive statistics are computed for baseline demographic, clinical and laboratory features according to etiology. Frequency tables were generated to reflect the etiology of cataract and all the values are determined at 5% significance level. Contingency tables were analyzed through chisquared statistics. Univariate type of parametric type of statistical analysis has been used.


**Table No.14** Type of Surgery done in the current sample of subjects

Type of surgery	n	Percent
Only ECCE	2	4.17
SICS with PC IOL	29	60.75
ECCE with PC IOL & PPC	1	2.08
ECCE with ACIOL	1	2.08
Phaco with PC IOL	15	30.92
Total	48	100



Various types of operations done on the subjects of the current study are shown in table No. 14 and figure No.11. Among these surgeries majority 60.75% is SICS with PC IOL. Next commonly done operation 30.92% is Phaco with PC IOL. In

a 4.17% of the cases only ECCE without any IOL implantation was done. ECCE with PC IOL and PPC has been tried in 2.08% of the cases. ECCE with ACIOL has been placed in 2.08% of the cases.



**Figure No.11** Type of Surgery done in the given Sample

### Limitations of the study

1. The sample size is small
2. Only cases attending Hospital were recruited for the study.

### Bibliography

1. Use of the Delphi process in paediatric cataract management Massimiliano Serafino<sup>1</sup>, Rupal H Trivedi<sup>2</sup>, Alex V Levin<sup>3</sup>, M Edward Wilson<sup>2</sup>, Paolo Nucci<sup>1</sup>, Scott R Lambert<sup>4</sup>, Ken K Nischal<sup>5</sup>, David A Plager<sup>6</sup>, Dominique Bremond-Gignac<sup>7</sup>, Ramesh Kekunnaya<sup>8</sup>, Sachiko Nishina<sup>9</sup>, Nasrin N Tehrani<sup>10</sup>, Marcelo C Ventura<sup>11</sup> Bjo.Bmj.com-2016
2. Short-term outcomes of dry pars plana posterior capsulotomy and anterior vitrectomy in paediatric cataract surgery using 25-gauge instruments, Y Huang, L XieBjo.Bmj.com-2010
3. Timing and approaches in congenital cataract surgery: a randomised controlled trial Prof Weirong Chen MDA† Erping Long MDA† Jingjing Chen MDA† Zhenzhen Liu PhDa† Zhuoling Lin MDA Qianzhong Cao PhDa Xiaohang Wu MDA Qiwei Wang PhDa Duoru Lin MDA Xiaoyan LiMDA Prof Lixia Luo PhDa BoQu PhDa Prof Haotian Lin PhDa Prof Yizhi Liu PhDa- The Lancet-October,2016-Elsevier
4. Comparing the astigmatic outcome after paediatric cataract surgery with different incisions Amit Gupta<sup>1</sup>, Muralidhar Ramappa<sup>2</sup>, Ramesh Kekunnaya<sup>1</sup>, Pravin K Vaddavalli<sup>2</sup>, Debajit Ray<sup>1</sup>, Sunita Chaurasia<sup>2</sup>, Harsha L Rao<sup>3</sup>,Chandra Sekar Garudadri<sup>3</sup> –Bjo.Bmj.com-2012
5. Outcome of paediatric cataract surgery with primary posterior capsulotomy and anterior vitrectomy using intra- operative preservative- free triamcinolone acetonide Rohit Gupta ,Jagat Ram ,JaspreetSukhija – Acta ophthalmologica-2014
6. Improvement for dry pars plana posterior capsulotomy and anterior vitrectomy using a 25-gauge vitrectomy system in paediatric cataract surgery Yunhai Dai<sup>1,2</sup>, Yusen

Huang<sup>2</sup>, Xiaoming Wu<sup>2</sup>, Lixin Xie<sup>2</sup>-  
British journal of Ophthalmology-2013-  
Bjo.Bmj.com

7. Long- term results of bilateral congenital cataract treated with early cataract surgery, aphakic glasses and secondary IOL implantation Dong- Hyun Kim ,Jeong Hun Kim, Seong- Joon Kim –Acta Ophthalmologica-2012
8. Precision pulse capsulotomy: an automated alternative to manual capsulorhexis in paediatric cataract Pratik Chougule<sup>1</sup>, Vivekanand Warkad<sup>2</sup>, Akshay Badakere<sup>1</sup>Ramesh Kekunnaya<sup>1</sup>- BMJ open Ophthalmology-2019.