



Inhalation of Vapor with Medication (Diclofenac Sodium, Menthol, Methyl Salicylate and N-Acetyl Cysteine) Reduces Oxygen Need and Hospital Stay in COVID-19 Patients – A Case Control Study

Authors

HN Sarker¹, Masum Ahmed², Linkon Dutta³

¹Professor (Ex), Medicine, Sher-E-Bangla Medical College, Barishal, Bangladesh

²Assistant professor, Respiratory Medicine, Sher-E-Bangla Medical College, Barishal, Bangladesh

³Senior staff nurse, Sher-E-Bangla Medical College Hospital, Barishal, Bangladesh

*Corresponding Author

Prof. H.N. Sarker

Abstract

Background: In the midst of the devastating COVID pandemic where there is no specific and effective treatment, traditional therapy may help to ease the patient's suffering. Inhalation of vapor (VP) is an essential home remedy for stuffy, running nose in common cold, influenza and sinusitis. Steam inhalation is helpful in destroying the capsid of the SARS-CoV-2 envelope and preventing infection. Vapor with diclofenac sodium, menthol, methyl salicylate and N-acetyl cysteine may augment this effect.

Objective: To evaluate the effect of inhalation of vapor with medication and to compare with inhalation of vapor without medication.

Methods and Materials: A case control study taken place in Corona unit, Sher-E-Bangla Medical College Hospital, Barishal. 43 patients with mild to moderate COVID-19 were participated in this study. All are RT-PCR positive cases. Among them 16 patients were in control group and 27 in study group. In study group they were given vapor with Diclofenac Sodium, Menthol, Methyl Salicylate and N-Acetyl Cysteine and control group they were given normal steam/aquatic vapor two times in a day.

Results: This study determined that after regular inhalation of vapor with above medication, oxygen saturation level increased in the study group 384.61% in the morning and 515.79% at night comparing the control group. Furthermore, patients of study group need to stay nearly 1 day less in hospital in comparison to control group.

Conclusion: Regular use of vapor with medication reduces oxygen need and hospital stay in COVID-19 patients compared to only steam vapor inhalation.

Keywords: Vapor, RT-PCR, oxygen saturation.

Introduction

The world is facing the second wave of Coronavirus Disease 2019 (COVID-19) pandemic which is the most troublesome challenge to public health. COVID-19 caused by severe acute

respiratory syndrome coronavirus-2 (SARS CoV-2) started in Hubei Province, People's Republic of China in December 2019^[1]. The virus has since spread globally rapidly and World Health Organization (WHO) declared COVID-19

pandemic on 11th March 2020. The second wave is running and nobody knows where we are in the course of this disease. It becomes a significant challenge for the public health, science, and medical sectors^[2].

SARS CoV-2 spreads by droplets from coughing, sneezing, talking and sighing. At present, it is fully recognised that the virus can spread by airborne^[3-5]. Although much has been discovered regarding the transmission and clinical feature, less is known about the pathophysiology of COVID-19. COVID-19 is primarily the disease of respiratory tract, at least initially. Disease pathophysiology of COVID-19 can be divided into three phases that correspond to different clinical stages of the disease^[6]. Stage 1: Asymptomatic state (initial 1–2 days of infection) - the inhaled virus SARS-CoV-2 likely binds to

epithelial cells in the nasal cavity through ACE2 receptors and starts replicating^[7-8]. Stage 2: Invasion and infection of the upper respiratory tract (next few days) - the virus propagates and migrates down the respiratory tract along the conducting airways. Due to the involvement of the upper airways, the disease manifests with symptoms of fever, fatigue, myalgia, runny nose, sneezing, cough, sore throat digestive symptoms (nausea, vomiting, abdominal pain, and diarrhea). Stage 3: Hypoxia, ground glass infiltrates, and progression to ARDS - Unfortunately, about 20% of the infected patients will progress to stage 3 disease and will develop pulmonary infiltrates and some of these will develop very severe disease (ARDS).

Clinical spectrum of COVID-19 disease is shown in Table 1.

Table 1 Clinical spectrum of COVID-19 disease^[9-10]

Severity of disease	Presentation
Asymptomatic	<ul style="list-style-type: none"> No clinical symptoms Positive nasal swab test Normal chest X-ray
Mild illness	<ul style="list-style-type: none"> Fever, sore throat, dry cough, malaise and body aches or Nausea, vomiting, abdominal pain, loose stools
Moderate illness	<ul style="list-style-type: none"> Symptoms of pneumonia (persistent fever and cough) without hypoxemia (SpO₂> 92%) Significant lesions on high-resolution CT chest
Severe illness	<ul style="list-style-type: none"> Pneumonia with hypoxemia (SpO₂<92%)
Critical state	<ul style="list-style-type: none"> Acute respiratory distress syndrome, along with shock, coagulation defects, encephalopathy, heart failure and acute kidney injury.

About 80% of the infected patients have mild disease and mostly restricted to the upper and conducting airways. These individuals may be monitored at home with conservative symptomatic therapy. Around 15% patients will have moderate disease and only 5% develop severe and critical illness.

Inhalation of vapor (VP) is an essential home remedy for stuffy, running nose in common cold, influenza and sinusitis. The biggest advantage of breathing in wet, warm steam is that it can help relieve nasal congestion and swelling of blood vessels. Moisture can also help to loose and thin

tenacious mucus in sinuses and airways, allowing them to empty more quickly. This allows the breathing to return to normal, if only for a brief time. Menthol is a substance which has a cooling and anesthetic effect. This also decreases the cough reflex and can soothe the dry throat.

For the casual viewer, steam inhalation is one of the most commonly used home remedies for soothing and opening the nasal passages and relieving the effects of a cold or sinus infection. According to the researchers, steam inhalation cycles are helpful in destroying the capsid of the SARS-CoV-2 envelope and preventing infection.

According to them, the European Pharmacopoeia VI edition has prescribed steam inhalations as a treatment for respiratory disorders^[11].

During last year when I and my other 4 family members had been suffering from COVID-19, we inhaled vapor to ease our dry cough. Initially, it gave temporary relief. For better relief I looked for menthol, but I got a gel that contained menthol in addition to methyl salicylate and diclofenac sodium and a nabule of 600 mg of N-acetyl cysteine near my hand. I mixed these two and put it in water in an electric tea-pot. We inhaled that medicated vapor 2-3 times daily during illness and gave better result to relieve cough, easy expectoration of sputum and an astonishing effect of improving oxygen saturation.

In fact, till now there is no specific and effective treatment for COVID-19. Where there is no specific and effective treatment, people and also we, the physicians look for the traditional remedy. Hypoxia is an integral and dangerous feature of severe and critical stage of COVID-19 and maintenance of oxygen saturation is the key to management.

The above observation lead me to conduct this study. In this study, we have tried to determine the effects of inhalation of vapor with medication and to compare with only vapor inhalation. As a consequence, we have taken two groups of population – one study group inhaling vapor with medication and another control group inhaling only vapor.

Method and Materials

Methods: Case control study.

Population: Adult patients with RT-PCR +ve mild to moderate COVID 19. Severe and critical disease patients had been excluded from the study.

Duration: One month

Study Place: Corona unit, Sher-E-Bangla Medical College Hospital, Barishal.

Sample Size: Total 43 patients (N). Among them study group (SG) had 27 respondents and control group (CG) had 16 respondents.

Materials: Suitable vapor producing electric teapot, medicine, pulse oximeter and a structured questionnaire.

Study Group: This group inhaled vapor containing Menthol 0.02%, Methyl salicylate 0.05%, N- Acetyl cysteine 1.2 gm%, and Diclofenac sodium 1gm% twice daily in addition to conventional treatment. That is, 100 gram of emulsion contain diclofenac sodium 1 gram, N-acetyl cysteine 1.2 gm, menthol 20 mg, and methyl salicylic acid 50 mg. These drugs have no systemic and local side effects in these small doses, though it may cause slight eye irritation.

Control group: This group inhaled plain aquatic vapor in addition to conventional treatment.

Data Collection Tools: Data has been collected for over a month from the study unit. The whole questionnaire has been administered by the researcher along with a research assistant. The researcher provided training and advocate the assistants for data collection including both subjective feelings (cough, chest pain, breathlessness) 10 minutes after inhalation and objective findings (pulse, respiratory rate and SpO₂) before and 10 minutes after inhalation and to see the accuracy of collected data. An excel file was prepared for daily data input.

Data Analysis: For data input, SPSS and Microsoft Excel software has been used. Moreover, for data analysis, SPSS version 23 has been used as principal software.

Data Quality Management: Data quality management was performed based on data integrity, completeness, validity, uniqueness, accuracy and consistency. Data quality management has been done by randomly checking data from the questionnaire and find out those errors like incomplete, invalid and irrelevant data from the data set. Data quality management team has been corrected those incomplete data as required.

Results

In this study 43 respondents participated and information was collect according to protocol. The

data were collected by the researchers and data were numerically coded and entered into using SPSS 23 version. This has been presented

graphically and thematically, and discussed based on the theme developed from the collected data.

Table-2: Distribution of age among study group and control group (*N=43; SG=27, CG=16*)

Age Range	Study Group	Control Group
	N (%)	N (%)
20-30 years	3 (11.1%)	0 (0.0%)
31-41 years	0 (0.0%)	2 (12.5%)
42-52 years	10 (37.0%)	5 (31.3%)
53-63 years	10 (37.0%)	4 (25.0%)
64-74 years	3 (11.1%)	3 (18.8%)
>75 years	1 (3.7%)	2 (12.5%)
Total	27 (100.0%)	16 (100%)

Table 2 showed that age of maximum patients are between 42-52 years in both study group (37.0%) and control group (31.3%).




Figure-1: Pie chart of age distribution of the patients.

Table-3: Distribution of gender among study group and control group (*N=43; SG=27, CG=16*)

Sex	Study Group	Control Group
	N (%)	N (%)
Male	18 (66.7%)	8 (50.0%)
Female	9 (33.3%)	8 (50.0%)

Table 3 showed that in study group most of the patients were male (66.7%) and in control group, male and females were equal in number (50%).

Table-4: Comparative table of average hospital stay between study group and control group (*N=43; SG=27, CG=16*)

Average Hospital Stay	Study group	Control Group	Difference of Hospital Stay in respect to control group
	7.41 days	8.25 days	0.84 \cong 1 day less

Above demonstrated table 4 showed that average hospital stay of study group and control group. Statistical analysis reported that those people who take vapor with medication (diclofenac sodium, menthol, methyl salicylate and N-acetyl cysteine) had to stay in hospital nearly one day less than control group.




Figure-2: Bar chart of duration of hospital stay.

Figure-2 shows that average hospital stay of study group is 1 day less than control group.

Table-5: Change of oxygen saturation in both study group and control group.

Description	Study Group Level of oxygen	Oxygen saturation changed in Study group	Control Group Level of oxygen	Oxygen saturation changed in control group	Difference of oxygen saturation changed with respect to control group	Significance
Average SPO ₂ – before VP in 8 AM	95.47	1	95.59	0.26	384.61% ↑	0.000*
Average SPO ₂ – 10 minute after VP from 8 AM	96.47		95.85			
Average SPO ₂ – before VP in 8 PM	95.74	0.98	95.70	0.19	515.79% ↑	0.000*
Average SPO ₂ – 10 minute after VP from 8 PM	96.72		95.89			
Total Average	96.1		95.75			

Table 5 shows that after taking vapor with medication helps to increase SpO₂ 384.61% and 515.79% respectively in morning and evening respectively in comparison to control group.




Figure-3: Correlation matrix of study group.

Figure- 3 shows that vaporization (with diclofenac sodium, menthol, methyl salicylate and n-acetyl cysteine) and oxygen saturation level are significantly associated.

Table-6: Table of paired sample T test among Study group.

Paired Samples Test		Paired Differences					t	df	Sig. (2-tailed)
		Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference				
					Lower	Upper			
Pair 1	Average SpO2-10 Min after VP from 8 AM - Average of SpO2-before VP in 8 AM	.99741	.36719	.07067	.85215	1.14266	14.114	26	.000
Pair 2	Average of SpO2-10 Min after VP in 8 PM - Average of SpO2-before VP in 8 PM	.97778	.32304	.06217	.84999	1.10557	15.728	26	.000

Table 6 reports the mean and standard deviation of the difference scores for each pair of variables Pair-1 (Mean .99741) and Pair-2 (Mean .97778). The mean is the difference between the sample means. The mean difference between Pair

1(Average SpO2-10 Min after VP from 8 AM - Average of SpO2-before VP in 8 AM) and Pair 2 (Average of SpO2-10 Min after VP in 8 PM - Average of SpO2-before VP in 8 PM) both are statistically significant at $\alpha = 0.05$. This is because

'Sig. (2-tailed)' or $p > 0.05$. In this study found that 95% confident that correlation of Average SpO₂-10 Min after VP from 8 AM - Average of SpO₂-before VP in 8 AM between 0.85215 to 1.14266 and Average of SpO₂-10 Min after VP in 8 PM - Average of SpO₂-before VP in 8 PM among .84999 to 1.10557 in the control group.

Discussion:

Steam inhalation can destroy the capsid of the SARS-CoV-2 envelope and is helpful in preventing COVID-19 infection^[11]. It also relieves congestion and loosen thick tenacious mucus making it easy to clear. Menthol decreases the cough reflex, can soothe the dry throat and helps to clear the airways. Methyl salicylate and diclofenac sodium are non-steroidal anti-inflammatory drugs, which may reduce pain and inflammation in the respiratory tract that is an important component of pathophysiology of COVID-19. N-acetyl cysteine is an established mucolytic agent which is usually used as an effervescent tablet or solution for nabulisation. Nabulisation is an aerosol generating procedure which is relatively contraindicated in COVID-19 patients.

Vapor inhalation is an essential home remedy in the initial stage of COVID 19 infection. Vapor with these medications loosen the thick mucus, helps to clear it easily, ease the breathing and speeds up removal of virus. In this study, we have tried to evaluate the effect of vapor with medications on mild to moderate COVID-19 patients and to compare with only vapor inhalation.

In this study, 43 patients were randomly selected. All these patients are RT-PCR positive mild to moderate COVID patients. They were divided into two groups –the study group and the control group. In table-2 and Figure-1, the age distribution of the patients among the two groups is shown. The maximum patients were between the age of 42-52 years in both study groups (37.0%) and the control group (31.3%).Table-3

shows that among all the respondents males were high in percentage in study group (66.7%).

In this study where patients of study group was given vapor with diclofenac sodium, menthol, methyl salicylate and N-acetyl cysteine at 8 AM and 8PM. On the other hand control group was given normal aquatic vapor at the same time. Maximum patients (87.50%) need to stay 5-10 days and 12.50% patients 1-5 days in the hospital. It was observed that patient of study group stayed in the hospital in an average 7.41 days. Besides, patients of control group were 8.25 days in hospital (Table-4). It shows that patients of study group stayed nearly 1 day less in hospital in comparison to control group (Fig-2). So it is easily understandable that regularvapor inhalation with medication helps to reduce hospital stay.

Pulse oximeter was used to measure oxygen level (SpO₂) of the participants. In the study group patients had average SpO₂95.47 before VP at 8 AM and96.47, 10 minute after VP at 8 AM, and average SpO₂ were 95.74before VP at 8 PM and 96.72,10 minute after VP at 8 PM. Oxygen saturation level was increased by 1 in the morning and 0.98 at night (Table-5).

On the other hand, in the control group, the average SpO₂ was 95.59 before VP at 8 AM and 95.85, 10 minute after VP at 8 AM, and the average SpO₂ was 95.70 before VP at 8 PM and 95.89,10 minutes after VP at 8 PM. Oxygen saturation level is increased by 0.26 in the morning and 0.19 at night (Table-5). It is clearly determined that oxygen saturation level is increased in the study group 384.61% in the morning and 515.79% at night comparing the control group after inhalation of vapor with medication. It is statistically significant at $\alpha=0.005$.

The mean and standard deviation of the difference scores for each pair of variables Pair-1 (Mean .99741) and Pair-2 (Mean .97778). The mean is the difference between the sample means. The mean difference between Pair 1(Average SpO₂-10 Min after VP from 8 AM - Average of SpO₂-before VP in 8 AM) and Pair 2 (Average of SpO₂-

10 Min after VP in 8 PM - Average of SpO₂-before VP in 8 PM) both are statistically significant at $\alpha = 0.05$. This is because 'Sig. (2-tailed)' or $p > 0.05$. In this study found that 95% confident that correlation of Average SpO₂-10 Min after VP from 8 AM - Average of SpO₂-before VP in 8 AM between 0.85215 to 1.14266 and Average of SpO₂-10 Min after VP in 8 PM - Average of SpO₂-before VP in 8 PM among .84999 to 1.10557 in the control group.

From the above discussion we can come to the point that vapor with diclofenac sodium, menthol, methyl salicylate and N-acetyl cysteine gives better result in COVID 19 patients in both reduction of oxygen needs and reduction of hospital stay compared with normal vapor inhalation.

Conclusion

In this situation of devastating COVID pandemic where there is no specific and effective treatment, traditional therapy may help to ease the patient's suffering. Regular vapor inhalation can prevent the deadly infection COVID 19 and vapor inhalation with above medication can help to reduce the symptoms, restore the breathing to normal, though only for a brief period, and reduce oxygen need and hospital stay.

This was a small study. So large studies are needed to verify the result.

References

1. WHO | Pneumonia of unknown cause – China. 2020 [cited 2021 Apr 24]; Available from: <https://www.who.int/csr/don/05-january-2020-pneumonia-of-unknown-cause-china/en/de>
2. Wit E, van Doremalen N, Falzarano D, Munster VJ. SARS and MERS: recent insights into emerging coronaviruses. *Nat Rev Microbiol*. 2016;14(8):523–34.
3. Asadi S, Wexler AS, Cappa CD, Barreda S, Bouvier NM, Ristenpart WD. Aerosol emission and superemission during human speech increase with voice loudness. *Sci Rep* 2019;9:2348. doi: 10.1038/s41598-019-38808-z pmid: 30787335
4. Scheuch G. Breathing is enough: for the spread of influenza virus and SARS-CoV-2 by breathing only. *J Aerosol Med Pulm Drug Deliv* 2020;33:230-4. doi: 10.1089/jamp.2020.1616 pmid: 32552296
5. Stadnytskyi V, Bax CE, Bax A, Anfinrud P. The airborne lifetime of small speech droplets and their potential importance in SARS-CoV-2 transmission. *Proc Natl AcadSci U S A* 2020;117:11875-7. doi: 10.1073/pnas.2006874117 pmid: 32404416
6. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72314 cases from the Chinese Center for Disease Control and Prevention. *JAMA* 2020.
7. Wan Y, Shang J, Graham R, et al. Receptor recognition by novel coronavirus from Wuhan: An analysis based on decade-long structural studies of SARS. *J Virol* 2020; 94: e00127-20.
8. Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* 2020.
9. Goyal P, Choi JJ, Pinheiro LC, et al. Clinical characteristics of COVID-19 in New York city. *N Engl J Med* 2020;382: 2372–4.
10. Guan WJ, Ni ZY, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020;382:1708–20.
11. Steam inhalation therapy found to inactivate SARS-CoV-2 virions [Internet]. *News-medical.net*. 2020 [cited 2021 Apr 24]. Available from: <https://www.news-medical.net/news/20201130/Steam-inhalation-therapy-found-to-inactivate-SARS-CoV-2-virions.aspx>.